

000007" 48208960





FIG. 1

TEMPERATURE RESPONSE OF IN VITRO r_1 FOR GdDTPA-BMA
ENCAPSULATED IN DPPC/DPPG LIPOSOMES (0.47T)

2 / 13

**FIG. 2**

TEMPERATURE RESPONSE OF MR SIGNAL INTENSITY FOR GdDTPA-BMA
ENCAPSULATED WITHIN DPPC/DPPG LIPOSOMES (2.0 T).




Figure 3. Gel phantom (a) containing inserts of DPPC/DPPE-based GdDTPA-BMA liposomes (labelled 3-10) and control glucose 5% solution (labelled 1-2); T_1 -w GRE images (2.0 T) of phantom prior to (b), after (c) 47 and (d) 63 minutes of radiofrequency heating, inhomogeneous signal intensity in gel is due to air bubbles; (e) difference image after subtraction of (b) from (d). Note that the signal intensity from inserts 3 and 5 is almost unchanged after heating as the temperature never exceeded T_c .




FIG.4




FIG.5

09680284-100600




FIG.6




FIG.7

00000T 49209960




FIG.8




FIG.9

[illegible]

[illegible]




FIG.12




FIG.13

09680284 100600




Figure 14




Figure 15




Figure 16

00900T"49208960




Figure 17




Figure 18




Figure 19




Figure 20

09680284-100600